

CONSTRAINING JET THEORIES USING NOVA OUTBURSTS

ALON RETTER¹
Draft version February 2, 2008

ABSTRACT

It is commonly accepted that jets have not been observed in cataclysmic variables (CVs) so far. This absence was recently explained by their low mass transfer rates compared with objects with jets. A mass accretion limit for jets in CVs was proposed to be $\dot{M} \sim 10^{-7} - 10^{-6} M_{\odot}/\text{yr}$. There was, however, a report of evidence for jets in V1494 Aql, the second nova that erupted in Aquila in 1999. In this work, we estimate the mass transfer rate of this system around the reported event and show that it is consistent with the above theoretical limit for jets. We further propose that the X-ray flare that was observed in this object may be connected with a jet as well. The appearance of jets in novae is actually expected since during the early decline from outburst some are alike supersoft X-ray sources, in which jets have been found. The detection of jets in novae also fits the suggestion that in addition to the presence of an accretion disk, a hot central source is required for the formation of jets. The observations of jets during the transition phase in V1494 Aql can be regarded as evidence for the early existence of the accretion disk in the system. This conclusion supports our previous suggestion for a link between the transition phase in novae and the re-establishment of the accretion disk. We further speculate that jets may be restricted to transient novae. If our ideas are confirmed, jets should be common in transient novae and may be formed several times during the transition phase and perhaps even long after it ends. In classical novae jets may be launched and observed in real time. As binary systems, these objects are easy to study. Novae may, therefore, be key systems in understanding the formation and evolution of jets and ideal targets to test and constrain jet theories. Adding novae to the classes of systems showing jets should be very exciting and rewarding.

Subject headings: stars: accretion, accretion disks—stars: novae—ISM: jets and outflows—stars: flare—stars: magnetic fields —stars: individual (V1494 Aql)

1. INTRODUCTION

Jets are one of the most spectacular astrophysical events. A jet is a narrow beam of matter or radiation emerging from an astronomical object. Usually two beams at opposite directions are observed from a central source. Collimated jets have been detected in young stellar objects, low mass and high mass X-ray binaries, supersoft X-ray sources, planetary nebula nuclei, symbiotic stars and active galactic nuclei.

Livio (2000) reviewed the observations and theory of jets and argued that the data of systems showing jets are consistent with the presence of accretion disks in all cases. Using a phenomenological approach, he also suggested that jets are launched from the center of the disk. It is commonly assumed that magnetic fields play a crucial role in the formation of jets and that they are accelerated and collimated hydromagnetically. Livio (2000) further proposed that the production of jets may require an additional heat or wind source probably associated with the central object. Yet, the jet mechanism is still obscure and so far numerous theories have been proposed to explain their formation.

Cataclysmic variables (CVs) contain a white dwarf star and a low mass red dwarf companion which fills its Roche lobe. Typically, the white dwarf accretes mass from an accretion disk that surrounds it. In most cases, jets have been associated with accretion disks, but it is widely accepted that jets have not been observed in CVs so far (e.g., Sokoloski et al. 2004; Hillwig, Livio & Honeycutt 2004). This latter perception is somewhat surprising, especially since jets have been detected in many symbiotic systems which contain accreting white dwarfs (Sokoloski et al. 2004; Galloway & Sokoloski 2004). It should be noted that an earlier report of jets in a CV (T Pyx) was based

on a misinterpretation of the data (O’Brien & Cohen 1998), so to date there has not been any convincing evidence of jets in CVs.

CVs can be divided into two main subgroups: dwarf novae and nova-like systems. Dwarf novae have relatively low mass transfer rates ($\dot{M} \sim 10^{-11} - 10^{-10} M_{\odot}/\text{yr}$) and their accretion disks are thermally unstable leading to dwarf nova outbursts typically every several months in their light curves. Nova-like systems have higher accretion rates ($\dot{M} \sim 10^{-9} - 10^{-8} M_{\odot}/\text{yr}$) resulting in stable accretion disks and no dwarf nova eruptions. After the white dwarf accretes enough material on its surface, and once its temperature and density reach some critical values, a thermo-nuclear-runaway occurs, giving rise to the observed nova outburst. It is believed that all CVs have nova eruptions every several ten thousands years (Warner 1995).

Soker & Regev (2003) pointed out recent observations that showed no difference in the X-ray properties of young stellar objects with or without jets (Getman et al. 2002). They, thus, argued that the observations are inconsistent with models that invoke the presence of magnetic fields for the launching mechanism of jets. Instead, they advanced a thermal model for the formation of jets in young stellar objects. It states that the accreted matter is shocked because of large gradients of physical quantities in the boundary layer near the accreting star. Soker & Lasota (2004) further applied this model to CVs to explain the absence of jets in these systems. They derived a rough estimate to the limiting mass transfer rate for jets in CVs: $\sim 10^{-7} - 10^{-6} M_{\odot}/\text{yr}$. As CVs typically accrete at much lower rates ($\dot{M} \lesssim 10^{-8} M_{\odot}/\text{yr}$ —e.g., Knigge & Livio 1998), they should not show jets.

¹ Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Lab, University Park, PA 16802-6305; retter@astro.psu.edu

1.1. The transition phase in classical novae

After the nova outburst, the white dwarf is quite hot and its cooling down process takes about a decade (Priyalnik 1986). The nova should have a high accretion rate that would gradually decrease in time. Classical novae may thus be excellent objects to check the evolution of jets if indeed they are formed in these systems. The accretion disk may be destroyed during the nova blast resulting in no accretion at all. However, strong evidence for the presence of the disk in novae several months-years after eruption have been found (Retter, Leibowitz & Ofek 1997; Skillman et al. 1997; see also Leibowitz et al. 1992; Retter, Leibowitz & Kovo-Kariti 1998). In addition, the white dwarf heated by the nova outburst may supply the extra energy source required for the formation of jets (Livio 2000). These systems may therefore, be ideal laboratories to test jet theories. The contribution from the white dwarf can be removed using theoretical models of its cooling (Priyalnik 1986; Somers & Naylor 1999). The observations or non-detection of jets at different stages of the nova decline can put strong constraints on jet models.

Classical novae can be broadly catalogued into three major subclasses according to the shape of their light curves. They either show a smooth decline, a deep minimum or oscillations during the transition phase several weeks-months after maximum light. The deep minimum in the light curve is explained by the formation of dust that may be restricted to systems with CO white dwarfs compared with ONeMg primaries in other novae (e.g., Rudy et al. 2003). The nature of the oscillations during the transition phase is, however, still under intensive debate.

Retter (2002) reviewed this issue and proposed a possible connection between the transition phase in classical novae and the re-formation of the accretion disk in a subclass of CVs called intermediate polars. These are systems whose inner part of the accretion disk is truncated by a moderate magnetic field of the white dwarf instead of a strong magnetic field, as in the case of polars. According to this idea, in a nova outburst it is easier to destroy the disk in intermediate polars as the inner part of the disk is missing. Retter (2002) thus proposed that the re-establishment of the accretion disk and the interaction between the inner part of the disk with the magnetic field of the white dwarf may form the oscillations observed during the transition phase. The test of this idea is very simple: novae with a transition phase (hereafter transient novae) should be intermediate polars and have fast spin periods. The detection of short-term periodicities in the X-ray light curves of two recent transient novae, V1494 Aql 1999 (Drake et al. 2003) and V4743 Sgr 2002 (Ness et al. 2003) is consistent with this suggestion. We note, however, that Drake et al. (2003) interpreted the variations in V1494 Aql as stellar pulsations of the hot white dwarf, while Ness et al. (2003) discussed both models as the cause for the X-ray fluctuations in V4743 Sgr.

In this work we investigate the first reliable report of jets in CVs, in V1494 Aql, which is a transient nova.

1.2. Nova V1494 Aql 1999

The outburst of V1494 Aql was discovered in 1999 December 1 (Pereira 1999). Reaching $V \sim 4$ at maximum a day later, it was one of the brightest novae in the century. Figure 1 presents the visual light curve of V1494 Aql during the first couple of years following discovery. A few months after the outburst the light curve was characterized by ~ 1 -2 mag oscillations with a quasi-period of 1-2 weeks typical of the transition phase in novae (Kiss & Thomson 2000; Iijima & Esenoglu 2003).

Extensive photometric observations of V1494 Aql revealed a periodic signal at 3.2 h (Retter et al. 2000; Bos et al. 2001; Pavlenko, Dudka & Baklanov 2003; Barsukova & Goranskii 2003; Kato et al. 2004). The folded light curve showed eclipses whose physical nature is still unknown. Bos et al. (2001) proposed that the accretion disk is eclipsed by the secondary star. Pavlenko et al. (2003) argued that the periodic variations can be explained by the presence of a strong magnetic field of the white dwarf and no accretion disk (i.e., polar model). Barsukova & Goranskii (2003) favored an intermediate polar model for the nova, with a truncated disk and a moderate magnetic field of the white dwarf. Recently, Hachisu, Kato & Kato (2004) demonstrated that the orbital light curve can be understood by the formation of a spiral structure in the accretion disk.

Iijima & Esenoglu (2003) analyzed spectroscopic observations of V1494 Aql during its early fading phase. Their data were intermittently taken during different stages of the decline from outburst, including the transition phase. Around one of the extrema of the long-term oscillations, two spectra showed symmetric emission features around the $H\alpha$ and $H\beta$ lines with a velocity of about 2800 km/s. This was interpreted by the authors as evidence for jets. In this work, we assess this suggestion and discuss its implications.

2. THE MASS TRANSFER RATE IN V1494 AQL

The presence of jets in V1494 Aql can be supported by estimating its mass accretion rate during the ‘jet event’ and showing that it is consistent with the theoretical limit. First, we assume that an accretion disk existed in this system in 2000 March. This assumption is justified by the fact that in most CVs the white dwarf accretes material from its companion star through a disk and strong evidence for its existence in novae several months-years after outburst has been detected (Section 1.1). In 2000 March, at the minimum of the oscillation, V1494 Aql was at $V \sim 10$ (Fig. 1), about 6 mags fainter than the main maximum of the outburst. Evidence for the presence of the accretion disk was first found in Nova V1974 Cyg 1992 when it was about 8 mags below maximum (Retter & Leibowitz 1998). The disk may have been present earlier in this system, although not as the dominant light source. Thus, it seems reasonable to assume that the accretion disk existed in V1494 Aql during the ‘jet event’. Conversely, we can claim that the detection of jets in V1494 Aql is an indication for the presence of the accretion disk in this nova in 2000 March.

The eclipses in V1494 Aql were discovered in 2000 June with an amplitude of 0.03 mag (Fig. 1; Retter et al. 2000). A year later, the eclipse depth was about 0.5 mag (Bos et al. 2001), and actually should be even larger after subtracting the contribution of light from the nearby background star (Kiss, Csák & Derekas 2004). Thus, in 2001 June the accretion disk was probably the main optical source.

The mass transfer rate of V1494 Aql will be estimated from its physical parameters. To simplify this calculation, we assume that the visual luminosity of V1494 Aql was completely dominated by the accretion disk in 2001 June and that the disk was not irradiated by the hot white dwarf. We follow Webbink et al. (1987) who developed a formula for the mass accretion rate of a CV as a function of its visual magnitude, distance, interstellar reddening and inclination angle. The visual magnitude of the nova in 2001 June was $V \sim 13.5$ (Fig. 1; Kiyota, Kato, & Yamaoka 2004). The distance to V1494 Aql and the interstellar extinction are taken as $d = 1.6$ kpc and $A_V = 2.0$ (Iijima &




FIG. 1.— The visual light curve of V1494 Aql during the first two years after discovery, taken from the VSNET (Variable Star Network). The nova had 1-2 mag oscillations with a quasi-period of 1-2 weeks during the transition phase, several months after peak outburst. Evidence for jets was found in 2000 March 16-17. V1494 Aql was classified as a supersoft X-ray source in 2000 August 6. An X-ray burst was observed in the system in 2000 October 1. Eclipses with a period of 3.2 h were first detected in the optical light curve of the nova in 2000 June with an amplitude of 0.03 mag. In 2001 June, the eclipse depth was much larger.

Esenoğlu 2003). The presence of eclipses constrains the inclination angle to high values, and it is assumed to be $i=80^\circ$. These values yield a mass transfer rate of about $5 \times 10^{-6} M_\odot/\text{yr}$.

The correct mass accretion rate of V1494 Aql in 2001 June should be somewhat lower than this value because of the assumption that the contribution from the white dwarf is negligible. The calculation shows, however, that if the accretion disk already existed at this time, the mass transfer rate was very high. If we further assume that during the ‘jet event’ in 2000 March, the mass accretion rate was of the same order (and it may have even been higher because the nova was brighter), then it obeys the critical limit for jets in CVs set by Soker & Lasota (2004).

Another independent way of estimating the accretion rate is to postulate that in the transition phase the accretion disk is destroyed during each oscillation, which we interpret as a mini-outburst (see below). An accretion disk in a typical dwarf nova before outburst has a mass of $\sim 10^{-9} M_\odot$ (e.g., Cannizzo, Gehrels & Mattei 2002). The mass of the disk in old novae and nova-like systems (as should be in V1494 Aql when it returns to quiescence) is probably even larger. Assuming that the disk is re-established between two successive mini-outbursts, thus within a few days, this is translated to mass transfer rates of the order of $10^{-7} M_\odot/\text{yr}$. This value should be regarded as a low limit to the mass accretion rate of the nova in 2000 March.

As a summary of this section, we conclude that if an accretion disk existed in V1494 Aql in 2000 March, then its mass transfer rate was probably in the range 10^{-7} – $5 \times 10^{-6} M_\odot/\text{yr}$, say $\sim 10^{-6} M_\odot/\text{yr}$.

3. DISCUSSION

We feel secure to argue that V1494 Aql had an accretion disk during the jet event and that the mass transfer rate at that time was relatively very high and almost certainly above the theoretical limit for jets despite the uncertainties in the calculations made in the previous section. To support this claim, we note that some novae have a supersoft X-ray stage for several months-years during the early decline from outburst (e.g., Orio, Covington & Ögelman 2001; Ness et al. 2003). Supersoft X-ray sources have high accretion rates and jets have been

observed in a few systems (see references in Livio 2000; Soker & Lasota 2004). V1494 Aql was classified as a supersoft X-ray source in 2000 August 6 (Drake et al. 2003). Therefore, the presence of high mass transfer rates and jets in this nova during its early fading phase is consistent with the observations of these systems as well.

On 2000 October 1, the X-ray emission from V1494 Aql increased suddenly by a factor of ~ 10 for about 15 minutes (Drake et al. 2003). This phenomenon is still unexplained. Our first estimate of the mass transfer rate in V1494 Aql ($\dot{M} \sim 5 \times 10^{-6} M_\odot/\text{yr}$) is for 2001 June, eight months after this event. In 2000 October, the mass accretion rate should have, therefore, been pretty large. We thus propose that this flare or burst might be connected with the formation of a jet as a result of a high mass transfer event.

Recent optical observations of Nova V4745 Sgr 2003 showed that spectra of this transient nova during the major maximum of its eruption and other maxima of the oscillations during the transition phase were alike. This discovery invoked an explanation of several repeating thermo-nuclear mini-outbursts as the cause of the oscillations (Csák et al. 2004). We note that these observations seem inconsistent with the explanation of the transition phase by winds (Shaviv 2001). Therefore, we refine Retter’s (2002) model (Section 1.1) and consider the following scenario for the transition phase: the nova outburst destroys the accretion disk in intermediate polars. After the white dwarf cools down and once its radiation pressure falls below a certain value, the disk is re-established within a few days, which leads to instantaneous high mass transfer rate. This triggers another mini nova outburst, that should demolish the disk again. The cycle may repeat several times with decreasing amplitudes until the mass transfer rate and the white dwarf temperature are too low to trigger another mini-outburst.

We suggest then that jets should be present during the transition phase in classical novae when presumably the accretion disk is re-established and the mass accretion rate is high enough, above the theoretical limit. They should be formed several times in accord with the number of mini-outbursts. In addition, if the mass transfer rate is still large enough, above

the critical value, collimated jets may be present even after the transition phase ends. The formation of jets should be ceased once the mass transfer rate drops below the critical limit. When the nova returns to quiescence, it should behave like a typical CV and not show jets.

The formation of jets has been linked with the presence of magnetic fields of the accretion disk itself or of the accreting star. Following our suggestion that the transition phase occurs in intermediate polars and assuming that jets are restricted to transient novae, we propose that the requirements for jets are the presence of an accretion disk and a moderate magnetic field of the inner source rather than the disk. This suggestion will be tested by future observations.

The thermal launching model presented by Soker & Lasota (2004) requires that the accretion disk be strongly disturbed at its inner part. This can be achieved by a boundary layer with a star, or by a strong stellar magnetic field (see second sentence in section 1.3 of Soker & Regev 2003). Therefore, the discussion in the present paper is valid for jets launched by systems where the disk inner radius is at the stellar surface as well as for jets from intermediate polars, where the disks are truncated by a moderate white dwarf magnetic field.

4. SUMMARY AND CONCLUSIONS

We find that during its jet event V1494 Aql had a high mass accretion rate, $\sim 10^{-6} M_{\odot}/\text{yr}$, consistent with the theoretical limit for jets in CVs (Soker & Lasota 2004). It is argued further that the peculiarities of V1494 Aql and V4743 Sgr in the X-ray may be explained by the ideas presented in this paper. According to our scenario when the nova decays and the accretion disk is reformed, jets should appear as a consequence of the temporary high accretion rate. The presence of jets in novae shortly after eruption seems natural because novae typically have a su-

persoft X-ray phase during the decline from outburst and jets have been detected in supersoft X-ray sources.

The evidence for jets in classical novae requires further support and it may be a coincidence that they appeared in a transient nova, nevertheless we speculate that jets may be restricted to transient novae. Furthermore, we propose that this phenomenon may be connected with the re-formation of the accretion disk and thus may be repeated several times before the mass transfer rate drops below the critical limit for jets.

We suggest that novae are ideal targets for the study of jets. This is because there is an evolution from high to low accretion rates within a few months or years in these systems in addition to a gradual change of the dominant light source from the hot white dwarf to the accretion disk. Further observations or non-detection of jets at different stages of the nova decline can put strong constraints on jet models. This paper thus calls for extensive X-ray, radio and optical spectroscopic monitoring of classical novae (and especially of transient novae) during their early decline stage. In addition, we encourage researchers to look for evidence of jets in archive high resolution spectroscopic data of novae.

The addition of classical novae to the known groups of objects showing jets should be quite exciting and may yield new insights on the formation of jets. It is very likely that novae, which are easy to study as binary systems, will supply the missing information on the jet mechanism.

We warmly thank M.T. Richards for many useful comments on an early draft of the paper and N. Soker for his encouragement to publish this work and for helpful discussions. This work was partially supported by a postdoctoral fellowship from Penn State University.

REFERENCES

- Barsukova, E.A., & Goranskii, V.P. 2003, *AstL*, 29, 195
 Bos, M., Retter, A., Cook, L., & Novak, R. 2001, *IAUC*, 7665
 Cannizzo, J.K., Gehrels, N., & Mattei, J.A. 2002, *ApJ*, 579, 760
 Csák, B., Kiss, L., Retter, A., Jacob, A., & Kaspi, S. 2004, *A&A*, accepted
 Drake, J.J. et al. 2003, *ApJ*, 584, 448
 Galloway, D.K., & Sokoloski, J.L., 2004, *ApJ*, submitted (astro-ph/0405436).
 Getman, K.V., Feigelson, E.D., Townsley, L., Bally, J., Lada, C.J., & Reipurth, B. 2002, *ApJ*, 575, 354
 Hachisu, I., Kato, M., & Kato, T. 2004, *ApJ*, 606, L139
 Hillwig, T., Livio, M., & Honeycutt, R.K. 2004, *PASP*, 116, 397
 Iijima, T., & Esenoglu, H.H. 2003, *A&A*, 404, 997
 Kato, T., Ishioka, R., Uemura, M., Starkey, D.R., & Krajci, T. 2004, *PASJ*, 56, 125
 Kiss, L.L., & Thomson, J.R. 2000, *A&A*, 355, L9
 Kiss, L.L., Csák, B., & Derekas, A. 2004, *A&A*, 416, 319
 Kiyota, S., Kato, T., & Yamaoka, H. 2004, *PASJ*, 56S, 193
 Knigge, C., & Livio, M. 1998, *MNRAS*, 297, 1079
 Leibowitz, E.M., Mendelson, H., Mashal, E., Prialnik, D., & Seitter, W.C. 1992, *ApJ*, 385, L49
 Livio, M. 2000, in *Cosmic Explosions: Tenth Astrophysics Conference*. AIP Conf. Proc., ed. S.S. Holt, & W.W. Zhang, 522, 275
 Ness, J.U. et al. 2003, *ApJ*, 594, L127
 O'Brien, T.J., & Cohen, J.G. 1998, *ApJ*, 498, L59
 Orio, M., Covington, J., & Ögelman, H. 2001, *A&A*, 373, 542
 Pavlenko, E.P., Dudka, O.I., & Baklanov, A.V. 2003, *Ap*, 46, 191
 Pereira, A. 1999, *IAUC*, 7323
 Prialnik, D. 1986, *ApJ*, 310, 222
 Retter, A. 2002, *AIP Conf. Ser.*, Vol 637, "Classical Nova explosions", eds. Hernanz M., José J., American Inst. of Physics, 279.
 Retter, A., & Leibowitz, E.M. 1998, *MNRAS*, 296, L37
 Retter, A., Leibowitz, E.M., & Ofek, E.O. 1997, *MNRAS*, 286, 745
 Retter, A., Leibowitz, E.M., & Kovo-Kariti, O. 1998, *MNRAS*, 293, 145
 Retter, A., Cook, L., Novak, R., Saxton, J.M., Jensen, L.T., Korcakova, D., & Janik, J. 2000, *IAUC*, 7537
 Rudy, R.J., Dimpfl, W.L., Lynch, D.K., Mazuk, S., Venturini, C.C., Wilson, J.C., Puetter, R.C., & Perry, R.B. 2003, *ApJ*, 596, 1229
 Shaviv, N. 2001, *MNRAS*, 326, 126
 Skillman, D.R., Harvey, D., Patterson, J., & Vanmunster, T. 1997, *PASP*, 109, 114
 Soker, N., & Regev, O. 2003, *A&A*, 406, 603
 Soker, N., & Lasota, J.P. 2004, *A&A*, 422, 1039
 Sokoloski, J.L., Kenyon, S.J., Brocksopp, C., Kaiser, C.R., & Kellogg, E.M., 2004, *proc. of IAU Colloq. 194, Compact Binaries in the Galaxy and Beyond*
 Somers, M.W., & Naylor, T. 1999, *A&A*, 352, 563
 Warner, B. 1995, *Cataclysmic Variable Stars*, Cambridge University Press, Cambridge
 Webbink, R.F., Livio, M., Truran, J.W., & Orio, M. 1987, *ApJ*, 314, 653